

# BUREAU OF WATER

South Carolina Department of Health and Environmental Control

## SHELLFISH MANAGEMENT AREA 2

---

### 2004 ANNUAL UPDATE

#### **Shellfish Sanitation Program**

Water Monitoring, Assessment and Protection Division  
Environmental Quality Control - Bureau of Water  
2600 Bull Street  
Columbia, South Carolina 29201

**July 2004**



**WEB ADDRESS:**

<http://www.scdhec.net/water/html/shellfish.html#reports>

# 2004 ANNUAL UPDATE

[ Data Through December 2003 ]

## Shellfish Management Area 2 Shellfish Sanitation Program



**Preparer:** Fred K. Earnhardt, District Shellfish Program Manager  
Waccamaw Environmental Quality Control District  
1705 Oak Street Plaza  
Myrtle Beach, South Carolina 29577

**Reviewers/Editors:**  
David G. Baize, Division Director  
(and) Charles Newell, Shellfish Program Manager  
Water Monitoring, Assessment, and Protection Division  
Environmental Quality Control - Bureau of Water  
2600 Bull Street  
Columbia, South Carolina 29201

*David G. Baize, Division Director*  
Water Monitoring, Assessment, and Protection Division  
Environmental Quality Control - Bureau of Water

## TABLE OF CONTENTS

### Shellfish Management Area 02 Annual Update

Summary .....	2
Introduction .....	2
Pollution Source Survey .....	5
Survey Procedures .....	5
Point Source Pollution .....	6
A. Municipal and Community Waste Treatment Facilities .....	6
B. Industrial Waste.....	6
C. Marinas .....	6
D. Radionuclides .....	6
Nonpoint Source Pollution.....	6
A. Urban and Suburban Stormwater Runoff.....	6
B. Agricultural Runoff.....	6
C. Individual Sewage Treatment and Disposal Systems .....	6
D. Wildlife and Domestic Animals.....	6
E. Boat Traffic .....	7
F. Marine Biotoxins .....	7
Hydrographic and Meteorological Characteristics .....	7
Water Quality Studies	8
Conclusions .....	9
Recommendations .....	9

## Figures and Tables

### Figures:

(1) Shellfish Harvesting Classification Prior to this Survey .....	12
(2) Current Shellfish Harvesting Classification & Sampling Stations .....	13
(3) Potential Pollution Sources.....	14

### Tables:

(1) Shellfish Water Quality Sampling Stations Description .....	11
(2) Fecal Coliform Bacteriological Data Summary Sheet (January 01, 2001 - December 31, 2003) .....	15
(3) Water Quality Sampling Stations Data .....	16
(4) Rainfall Data (A: 2001 / B: 2002/ C: 2003) .....	17

**ANNUAL UPDATE**  
**Shellfish Management Area 02**  
**SCDHEC EQC Bureau of Water**

**Data Inclusive Dates:**

01 / 01 / 01 thru 12 / 31 / 03

**Classification Change:**

   Yes   X   No

**Shoreline Survey Completed:**   Yes  

**(I)ncreased/(D)ecreased/(N)one:**

  N   Approved

  N   Cond. Approved

  N   Restricted

  N   Prohibited

**Prior Report & Date:** Annual-2003

**SUMMARY**

Area 02 is primarily affected by nonpoint source pollution due to dense development. The shoreline reconnaissance and bacteriological data review indicate that the present restricted classification is appropriate. No depuration harvest permits should be issued for this area.

**INTRODUCTION**

**PURPOSE AND SCOPE**

The authority to regulate the harvest, sanitation, processing and handling of shellfish is granted to the South Carolina Department of Health and Environmental Control by Section 44-1-140 of the Code of Laws of South Carolina, 1976, as amended. The Department promulgated Regulation 61-47, which provides the rules used to implement this authority and outlines the requirements applied in regulating shellfish sanitation in the State. This regulation specifically addresses classification of shellfish harvesting areas and requires that all areas be examined by sanitary and bacteriological surveys and classified into an appropriate shellfish harvesting classification.

The National Shellfish Sanitation Program (NSSP) Guide For The Control Of Molluscan Shellfish is used by the United States Food and Drug Administration (USFDA) to evaluate state shellfish sanitation programs. The NSSP Model Ordinance requires that a sanitary survey be in place for each growing area prior to its use as a source of shellfish for human consumption and prior to the area's classification as Approved, Conditionally Approved, Restricted, or Conditionally Restricted. Each sanitary survey shall be updated on an annual basis and accurately reflect changes which have occurred within the area. Requirement of the annual reevaluation include, at a minimum, field observations of pollution sources, an analysis of water quality data consisting of the past year's data in combination with appropriate previously collected data, review of reports and effluent samples from pollution sources, and review of performance standards for discharges impacting the growing area. A brief report

documenting the findings shall also be provided.

The following criteria consistent with the NSSP Model Ordinance and S. C. Regulation 61-47 are used in establishing shellfish harvesting classifications:

**Approved** - Growing areas shall be classified Approved when the sanitary survey concludes that fecal material, pathogenic microorganisms, and poisonous or deleterious substances are not present in concentrations which would render shellfish unsafe for human consumption. The Approved area classification shall be designated based upon a sanitary survey which includes water samples collected from stations in the designated area adjacent to actual or potential sources of pollution. For waters sampled under adverse pollution conditions, the median fecal coliform Most Probable Number (MPN) or the geometric mean MPN shall not exceed fourteen per one hundred milliliters, and not more than ten percent of the samples shall exceed a fecal coliform MPN of forty-three per one hundred milliliters (per five tube decimal dilution). For waters sampled under a systematic random sampling plan, the geometric mean fecal coliform Most Probable Number (MPN) shall not exceed fourteen per one hundred milliliters, and the estimated ninetieth percentile shall not exceed an MPN of forty three (per five tube decimal dilution). Computation of the estimated ninetieth percentile shall be obtained using NSSP Guidelines.

**Conditionally Approved** - Growing areas may be classified Conditionally Approved when they are subject to temporary conditions of actual or potential pollution. When such events are predictable, as in the malfunction of wastewater treatment facilities, non-point source pollution from rainfall runoff, discharge of a major river, or potential discharges from dock or harbor facilities that may affect water quality, a management plan describing conditions under which harvesting will be allowed shall be adopted by the Department prior to classifying an area as Conditionally Approved. Where appropriate, the management plan for each Conditionally Approved area shall include performance standards for sources of controllable pollution, e.g., wastewater treatment and collection systems, evaluation of each source of pollution, and means of rapidly closing and subsequent reopening areas to shellfish harvesting. Memorandums of agreements shall be a part of these management plans where appropriate.

**Restricted** - Growing areas shall be classified Restricted when sanitary survey data show a limited degree of pollution or the presence of deleterious or poisonous substances to a degree which may cause the water quality to fluctuate unpredictably or at such a frequency that a Conditionally Approved classification is not feasible. Shellfish may be harvested from areas classified as Restricted only for the purposes of relaying or depuration and only by special permit issued by the Department and under Department supervision.

For Restricted areas to be utilized as a source of shellstock for depuration, or as source water for depuration, the fecal coliform geometric mean MPN of restricted waters sampled under adverse pollution conditions shall not exceed eighty-eight per one hundred milliliters and not more than ten percent of the samples shall exceed a MPN of two hundred and sixty per one hundred milliliters for a five tube decimal dilution test. For waters sampled under a systematic random sampling plan, the fecal

coliform geometric mean MPN shall not exceed eighty-eight per one hundred milliliters and the estimated ninetieth percentile shall not exceed an MPN of two hundred and sixty (five tube decimal dilution). Computation of the estimated ninetieth percentile shall be obtained using NSSP guidelines.

**Conditionally Restricted** - Growing areas may be classified Conditionally Restricted when they are subject to temporary conditions of actual or potential pollution. When such events are predictable, as in the malfunction of wastewater treatment facilities, non-point source pollution from rainfall runoff, discharge of a major river, or potential discharges from dock or harbor facilities that may affect water quality, a management plan describing conditions under which harvesting will be allowed shall be prepared by the Department prior to classifying an area as Conditionally Restricted. Where appropriate, the management plan for each Conditionally Restricted area shall include performance standards for sources of controllable pollution (e.g., wastewater treatment and collection systems and an evaluation of each source of pollution) and description of the means of rapidly closing and subsequent reopening areas to shellfish harvesting. Memorandums of agreements shall be a part of these management plans where appropriate. Shellfish may be harvested from areas classified as Conditionally Restricted only for the purposes of relaying or depuration and only by permit issued by the Department and under Department supervision. For Conditionally Restricted areas to be utilized as a source of shellstock for depuration, the fecal coliform geometric mean MPN of Conditionally Restricted waters sampled under adverse pollution conditions shall not exceed eighty-eight per one hundred milliliters and not more than ten percent of the samples shall exceed a MPN of two hundred and sixty per one hundred milliliters for a five tube decimal dilution test. For waters sampled under a systematic random sampling plan, the fecal coliform geometric mean MPN shall not exceed eighty-eight per one hundred milliliters and the estimated ninetieth percentile shall not exceed an MPN of two hundred and sixty (five tube decimal dilution). Computation of the estimated ninetieth percentile shall be obtained using NSSP guidelines.

**Prohibited** - Growing areas are classified Prohibited if there is no current sanitary survey or if the sanitary survey or monitoring data show unsafe levels of fecal material, pathogenic microorganisms, or poisonous or deleterious substances in the growing area or indicate that such substances could potentially reach quantities which could render shellfish unfit or unsafe for human consumption.

## **BACKGROUND INFORMATION**

Area 02 consists of three separate estuaries. White Point Swash, located in the Windy Hill section of North Myrtle Beach, is bordered to the northwest by U. S. Highway 17, to the northeast by mainland residential portions of Windy Hill, North Myrtle Beach, to the southwest by the residential development of Briarcliff Acres, and to the southeast by the Atlantic Ocean. Singleton Swash is located approximately six miles south of White Point Swash and is bordered generally to the west by U. S. Highway 17, to the south by the Dunes Golf and Beach Club of Myrtle Beach, to the north by Lake Arrowhead Road, and on the east by Shore Drive and the Atlantic Ocean. Cane Patch Swash is located within the Myrtle Beach city limits. It is bordered by U. S. Highway 17 on the west, on the north by 68th Avenue North, on the south by 66th Avenue North, and on the east by the Atlantic Ocean. (Figure 1) The combined area of shellfish waters in Area 02 is approximately 100 acres.

Although the area is small, a substantial shellfish resource does exist, necessitating review and classification.

The harvesting classification of Area 02 prior to this sanitary survey was as follows:

Prohibited: None;

Restricted     1) All waters of White Point Swash;  
                    2) All waters of Singleton Swash;  
                    3) All waters of Cane Patch Swash;

Conditionally Approved: None;

Approved: None.

The shellfish industry in South Carolina is based primarily on the harvest of the eastern oyster (*Crassostrea virginica*) and hard clams, which include both the northern clam (*Mercenaria mercenaria*) and several small populations of the southern clam (*Mercenaria campechiensis*). The South Carolina Department of Health and Environmental Control currently disallows harvesting of oysters and clams within Area 02 for direct marketing purposes. No relay projects have been permitted during the past three year review period.

Shellfish harvesting season in South Carolina normally extends from September 16 through May 14. The South Carolina Department of Natural Resources (SCDNR) has the authority to alter the shellfish harvesting season for resource management purposes and grant permits for year-round mariculture operations. Additionally, the South Carolina Department of Health and Environmental Control has the authority to prohibit shellfish harvesting when necessary to ensure that shellfish harvested in South Carolina waters are safe for human consumption.

## **POLLUTION SOURCE SURVEY**

### **SURVEY PROCEDURES**

Shoreline surveys of Area 02 were conducted by the Waccamaw District Shellfish Sanitation staff during the survey period and are ongoing. Extensive visual examination of lands adjacent to the waters of Area 02 was conducted in order to determine type and density of structures, location of significant concentrations of domestic animals, and other potential sources of pollution entering shellfish growing waters. Additionally, Waccamaw EQC District personnel, in conjunction with the Department's ocean water quality monitoring program, have extensively examined lands surrounding White Point Swash.

## POINT SOURCE POLLUTION

- A. **Municipal and Community Waste Treatment Facilities** - The majority of this area has central sewer and is serviced by the cities of Myrtle Beach and North Myrtle Beach. Discharge of effluent from their facilities is to the Atlantic Intracoastal Waterway and does not affect the shellfish growing waters in Area 02. The town of Briarcliff Acres, however, is primarily serviced by individual sewage treatment and disposal systems (ISTDs).
- B. **Industrial Waste** - There are no permitted point source discharges of industrial waste in Area 02.
- C. **Marinas** - There are no marinas located in Area 02 due to lack of navigable channels.
- D. **Radionuclides** - Sources of radionuclides have not been identified within Area 02, and radionuclide monitoring has not been conducted. No poisonous or deleterious substances have been identified.

## NONPOINT SOURCE POLLUTION

- A **Urban and Suburban Stormwater Runoff** - Nonpoint source runoff is the apparent primary contributor to elevated fecal coliform bacteria levels in the area. White Point Swash receives drainage from a densely developed portion of the City of North Myrtle Beach. Development includes an extensive residential area in addition to a family campground and numerous condominiums, restaurants, gift shops, and amusement facilities. Singleton Swash receives drainage from Singleton Lake and several small ponds within the management area. Development is extremely dense, primarily consisting of high-rise and smaller stick built condominiums, campgrounds, and golf courses. Cane Patch Swash serves as a drainage outlet for the City of Myrtle Beach. Drainage is from a series of small lakes located west of U. S. Highway 17. Development is dense with single and multi-family residences and numerous small businesses.
- B **Agricultural Runoff** - There are no commercial agricultural activities conducted adjacent to the waters of Area 02, and sampling for pesticides and herbicides has not been conducted.
- C **Individual Sewage Treatment and Disposal (ISTD) Systems** - Individual sewage treatment and disposal (ISTD) systems are known to exist in the North Myrtle Beach areas bordering White Point Swash and Briarcliff acres; however, exact numbers and distribution have not been identified.
- D **Wildlife and Domestic Animals** - Wildlife in Area 02 consists primarily of deer, raccoons, and opossums. These populations, in combination with domestic cats and dogs, are contributors to nonpoint source pollution. Effective resource management of deer populations coupled with a dramatic loss of natural habitat due to continued development within, and on



lands adjacent to, Area 02, has resulted in efforts to reduce the deer population within the town Briarcliff Acres. Briarcliff Acres is approximately one mile southwest of White Point Swash.

- E. Boat Traffic** - The lack of navigable channels within Area 02 generally precludes watercraft use.
- F. Marine Biotoxins** - During the winter and spring of 1988, South Carolina experienced an occurrence of "Red Tide", specifically *Ptychodiscus brevis* (*G. Breve*), which affected water quality in Area 01. There has been no documented reoccurrences of this organism at levels requiring emergency response in South Carolina waters subsequent to the 1988 event. The Department participates in a State Task Group on Toxic Algae and operates a toxic algae emergency response team.

## **HYDROGRAPHIC AND METEOROLOGIC CHARACTERISTICS**

### **PHYSIOGRAPHY**

White Point Swash is approximately 650 by 500 meters and includes approximately 42 acres of habitat suitable for shellfish production. Immediately adjacent to White Point Swash is a series of marsh areas that extend approximately 2000 meters in a northeast - southwesterly direction and parallel the beachfront. This area averages less than 100 meters in width. Total area is approximately 35 acres. A single shallow swash connects the White Point estuary with the Atlantic Ocean.

Singleton Swash contains approximately 41 acres of bottoms suitable for shellfish production. A single shallow swash immediately adjacent to the Dunes Golf and Beach Club connects the Singleton Swash estuary with the Atlantic Ocean.

Cane Patch Swash contains less than three acres of bottoms suitable for shellfish production. Connection to the Atlantic Ocean is via a shallow ditch and 36-inch drain culvert.

**Tides** - Tides along the beaches in Myrtle Beach/North Myrtle Beach are semidiurnal, consisting of two low and two high tides each lunar day. Mean tidal ranges are 5.00 - 5.06 feet during normal tides and 5.70 - 5.87 feet during spring tides (Tides and Currents for Windows, Version 2.2, Nautical Software Inc.).

**Precipitation** - Precipitation in Area 02 averages between 50 and 52 inches per year with the heaviest rainfall occurring during the summer months. Tropical storms and hurricanes occasionally produce extremely large amounts of rainfall.

During winter months rainfall is more uniform in nature; heavy, short-term rainfall events are uncommon- yet occasional intense thunderstorms associated with rapidly moving low pressure systems may generate heavy rains. Precipitation rarely occurs in the form of snow or ice, however, low pressure

systems infrequently cross Florida and become stationary just offshore of the Georgia coast. Such was the case in late December 1989. This storm system resulted in approximately 14 inches of snow along Horry County's beaches. Spring weather patterns are often extremely dynamic with associated thunderstorms and severe weather conditions.

**Winds** - Prevailing winds along the northern portion of the South Carolina coast are from the southwest during spring and south/southwest during the summer. During autumn wind direction is generally from the Northeast. Winter winds fluctuate between Northeast and Southwest. Wind speeds average less than 10 mph, however, strong weather systems may generate winds in excess of 25 mph. Tropical storms and hurricanes frequently occur.

**River Discharges** - There are no rivers in close proximity to Area 02. Freshwater input occurs via localized precipitation and resulting runoff.

**Currents** - Currents are tidally generated, although wind speed and direction may cause affect current velocities. Flow reverses direction approximately every six hours.

## **WATER QUALITY STUDIES**

### **DESCRIPTION OF THE PROGRAM**

The Department currently utilizes a systematic random sampling (SRS) strategy within Area 02 in lieu of sampling under adverse pollution conditions. In order to comply with NSSP guidelines, a minimum of thirty samples are required to be collected and analyzed from each station during the review period. Sampling dates are computer generated prior to the beginning of each quarterly period thereby insuring random selection with respect to tidal stage and weather. Day of week selection criteria is limited to Mondays, Tuesdays, and Wednesdays due to shipping requirements and laboratory manpower constraints. Sample schedules are rarely altered.

During July 1998, an updated data analysis procedure was formalized. Samples utilized for classification purposes are limited to those samples collected in accordance with the SRS for a 36-month period beginning January 1 and ending December 31. This allows for a maximum of 36 samples per station yet provides a six-sample 'cushion' (above the NSSP required 30 minimum) for broken samples, lab error, breakdowns, etc. This also allows each annual report to meet the NSSP Triennial Review sampling criteria.

One hundred and eight surface water quality samples (<1.0 ft. deep) were collected for bacteriological analyses and classification purposes from three active water quality sampling stations in Area 02 during the period 01/01/00 through 12/31/02. The samples were collected in 120 ml amber glass bottles, immediately placed on ice and transported by bus to the South Carolina Department of Health and Environmental Control's Trident District Environmental Quality Control laboratory at North Charleston, South Carolina. An additional 120 ml water sample was included with each shipment as a

temperature control. Upon receipt at the laboratory, sample sets that exceeded a 30-hour holding time or contained a temperature control >10 degrees C. were discarded. Samples collected after September 1, 1986 have been analyzed using the five tube/three dilution modified A-1 method described by Nuefeld (1985<sup>1</sup>).

Surface water temperatures were measured utilizing hand-held, laboratory-quality calibrated centigrade thermometers. Salinity measurements were measured in the laboratory using automatic temperature compensated refractometers. Additional field data include ambient air temperature, wind direction, tidal stage and date and time of sampling. Tidal stages were determined Nautical Software's Tides and Currents, Version 2.2 (1996).

## **MONITORING RESULTS**

Stations 02 and 03 exhibited fecal coliform geometric means in excess of 88 MPN/100 ml. All stations exceeded an estimated 90th percentile fecal coliform MPN/100 ml. value of 260.

## **CONCLUSIONS**

Based on review of fecal coliform bacteriological data summaries and the pollution source survey, Area 02 is impacted by one primary source of pollution. There are no specific point source discharges and the conclusion is that, primarily due to dense development, nonpoint source runoff is the prime contributor to elevated fecal coliform bacteria levels in the area.

## **RECOMMENDATIONS**

The shoreline reconnaissance and bacteriological data review of shellfish growing Area 02 indicated that the current restricted classification is appropriate. Due to the excessive fecal coliform geometric means and estimated 90th percentile values, no depuration activities should be allowed. The harvesting classification of Area 02 is recommended to remain as follows:

Prohibited:	None
Restricted:	1) All waters of White Point Swash; 2) All waters of Singleton Swash; and 3) All waters of Cane Patch Swash;
Conditionally Approved:	None

---

<sup>1</sup>Nuefeld, N. 1985. Procedures for the bacteriological examination of seawater and shellfish. In: A.E. Greenberg and D.A. Hunt (eds.) Laboratory procedures for the examination of seawater and shellfish, Fifth Edition. American Public Health Association, Washington, D.C. p. 37-63.

Approved:       None.

**Station Addition/Deactivation/Modification:**       None

**TABLE #1**

**Shellfish Management Area 02  
WATER QUALITY SAMPLING STATIONS DESCRIPTION**

<b><u>Station</u></b>	<b><u>Description</u></b>
01	White Point Swash
02	Singleton Swash
03	Canepatch Swash
(Total 3)	

Figure 1.  
Shellfish Management Area 02  
Prior Classification




Figure 2.  
Shellfish Management Area 02  
Current Classification





Figure 3.  
Shellfish Management Area 02  
Potential Pollution Sources





**TABLE #2**  
**Shellfish Management Area 02**

***FECAL COLIFORM BACTERIOLOGICAL DATA SUMMARY***  
***from Shellfish Water Quality Sampling Stations between***

**January 01, 2001 through December 31, 2003**

<b>Station ?</b>	<b>1</b>	<b>2</b>	<b>3</b>								
<b>Samples</b>	36	36	36								
<b>GeoMean</b>	83.9	104.9	468.3								
<b>90th %ile</b>	946	1347	3489								
<b>Water Qlty</b>	RND	RND	RND								
<b>Classification</b>	R	R	R								

<b>Station ?</b>											
<b>Samples</b>											
<b>GeoMean</b>											
<b>90th %ile</b>											
<b>Water Qlty</b>											
<b>Classification</b>											

<b>Station ?</b>											
<b>Samples</b>											
<b>GeoMean</b>											
<b>90th %ile</b>											
<b>Water Qlty</b>											
<b>Classification</b>											

**A** - Approved      **CA** - Conditionally Approved      **R** - Restricted  
**RND** - Restricted/No Depuration      **P** - Prohibited

**TABLE #3**

**WATER QUALITY SAMPLING STATIONS DATA**

---

**Shellfish Management Area 02**

**January 01, 2001 through December 31, 2003**

## **BACTERIOLOGICAL DATA**

Data for each shellfish station listed in this report's "Fecal Coliform Bacteriological Data Summary Table" and in other shellfish reports, can be obtained through South Carolina's Department of Health and Environmental Control - Freedom of Information office at the address below.

Freedom of Information  
2600 Bull Street  
Columbia, SC 29201

Any explanation or clarity needed on the report's content can be obtained by contacting the preparer(s), and/or reviewer(s) listed on the cover page.

**TABLE #4 (A-C)**

**RAINFALL DATA**

---

**Shellfish Management Area 02**

**SOURCE:**

*South Carolina Department of Natural Resources,  
Land, Water and Conservation Division,  
South Carolina State Climatology Office*

# ANNUAL TABLE OF RAINFALL DATA

*SOURCE: Department of Natural Resources*

*Land, Water and Conservation Division*

*State Climatology Office (Myrtle Beach, SC / Station #2)*

2001	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
1st	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.15	0.00	0.00	0.00
2nd	0.00	0.00	0.00	0.00	0.00	0.54	0.87	0.00	0.01	0.00	0.02	0.00
3rd	0.00	0.00	0.61	0.00	0.00	0.00	1.00	0.00	1.34	0.00	0.00	0.00
4th	0.00	0.60	1.24	0.05	0.00	1.02	1.10	0.00	1.65	0.00	0.00	0.00
5th	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.00	0.32	0.00	0.00	0.00
6th	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
7th	0.00	0.00	0.00	0.00	0.00	0.00	0.13	0.39	0.00	0.10	0.00	0.00
8th	0.00	0.00	0.00	0.00	0.00	1.10	0.00	0.00	0.00	0.00	0.00	0.00
9th	0.24	0.00	0.00	0.00	0.00	1.20	0.00	0.00	0.00	0.00	0.00	0.01
10th	0.00	0.00	0.03	0.00	0.00	0.01	0.46	0.00	0.00	0.00	0.00	0.01
11th	0.00	0.20	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
12th	0.03	0.52	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	2.20
13th	0.50	1.35	1.06	0.00	0.05	--	1.50	0.00	0.00	0.00	0.00	--
14th	0.00	0.00	0.00	0.10	0.00	0.46	0.28	0.35	0.00	0.00	0.00	0.00
15th	0.08	0.00	0.26	0.00	0.00	0.58	0.00	0.09	0.90	0.10	0.00	0.00
16th	0.00	0.00	1.24	0.00	0.00	0.24	0.00	0.00	0.00	0.06	0.00	0.00
17th	0.03	0.03	0.00	0.00	0.00	0.35	0.00	0.00	0.00	0.00	0.00	0.00
18th	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08
19th	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00
20th	0.03	0.00	0.05	0.00	0.00	0.00	0.66	0.14	0.00	0.00	0.00	0.00
21st	0.02	0.00	2.00	0.00	0.64	0.00	0.00	0.09	0.00	0.00	0.00	0.00
22th	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	1.31	0.00	0.00	0.00
23rd	0.00	0.00	0.00	0.00	0.36	0.10	0.36	0.00	0.00	0.00	0.00	0.00
24th	0.00	0.00	0.00	0.00	0.00	0.00	0.70	0.00	0.01	0.00	1.10	0.05
25th	0.00	0.00	0.00	0.72	0.00	0.40	0.01	0.04	0.05	0.00	0.66	0.00
26th	0.00	0.00	0.15	0.36	0.00	0.10	0.00	0.00	0.00	0.00	0.03	0.00
27th	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
28th	0.00	0.10	0.00	0.00	0.00	0.00	0.85	0.00	0.00	0.00	0.00	0.00
29th	0.00	--	0.40	0.00	0.05	0.00	0.00	0.00	0.00	0.00	--	0.00
30th	0.00	--	0.60	0.00	0.95	0.00	0.29	--	0.00	0.00	0.02	0.00
31st	0.10	--	0.00	0.00	0.00		0.00		--	0.00	--	0.00

(Monthly Figures below)

Year's Rainfall Total:

40.51

SUM	1.03	2.80	7.66	1.23	2.10	6.11	8.29	1.10	5.75	0.26	1.83	2.35
MAX	0.50	1.35	2.00	0.72	0.95	1.20	1.50	0.39	1.65	0.10	1.10	2.20
MIN	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
AVG	0.03	0.10	0.25	0.04	0.07	0.21	0.27	0.04	0.19	0.01	0.06	0.08

"ND"or"--" = missing data / Shaded Cells = date samples were taken  
(Shellfish Management Areas 02 and 03)

# ANNUAL TABLE OF RAINFALL DATA

*SOURCE: Department of Natural Resources*

*Land, Water and Conservation Division*

*State Climatology Office (Myrtle Beach, SC / Station #2)*

2002	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
1st	0.00	0.00	0.00	0.42	0.00	0.00	0.00	0.22	0.70	0.00	0.00	0.00
2nd	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00	0.00
3rd	0.49	0.00	1.30	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
4th	0.15	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00
5th	0.00	0.00	0.00	0.00	0.25	0.00	0.00	0.80	0.00	0.00	0.10	0.00
6th	0.26	0.00	0.00	0.00	0.00	0.00	0.66	0.00	0.10	0.00	0.00	0.40
7th	0.82	1.65	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
8th	0.00	0.30	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.00
9th	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00
10th	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.25
11th	0.00	0.30	0.00	0.03	0.90	0.00	0.00	0.00	0.00	0.35	0.10	0.97
12th	0.00	0.00	0.00	0.03	0.19	0.00	1.48	0.00	0.00	0.25	1.30	0.97
13th	0.38	0.00	0.52	0.00	0.00	0.00	0.01	0.00	0.00	0.00	1.35	0.30
14th	0.01	0.00	0.00	0.00	1.20	0.00	0.25	0.00	0.05	0.00	0.05	0.50
15th	0.50	0.00	0.00	0.00	0.00	0.30	0.24	2.20	0.40	0.30	0.00	0.00
16th	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.04	0.35	0.00	0.00
17th	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.50	0.00
18th	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.25	0.00	0.00	0.02	0.00
19th	0.00	0.00	0.00	0.00	0.50	0.35	0.00	0.28	0.15	0.00	0.00	0.00
20th	0.12	0.00	0.00	0.00	0.05	0.10	0.00	0.05	0.00	0.00	0.00	0.20
21st	0.00	0.07	0.00	0.00	0.00	0.27	2.30	0.00	0.00	0.00	0.00	0.00
22th	0.08	0.00	0.10	0.00	0.00	0.19	0.02	0.00	0.00	0.20	0.00	0.00
23rd	0.00	0.00	0.00	0.00	0.00	1.54	1.75	0.00	0.00	0.30	0.00	0.00
24th	0.05	0.00	0.00	0.00	0.00	0.05	0.06	0.00	0.96	0.05	0.00	1.20
25th	0.08	0.00	0.00	0.00	0.00	0.03	0.03	1.10	0.00	0.00	0.00	0.00
26th	0.00	0.00	0.00	0.36	0.00	0.00	0.00	0.23	1.50	0.00	0.00	0.00
27th	0.00	0.00	0.20	0.00	0.00	0.02	0.00	0.10	0.01	0.20	0.00	0.00
28th	0.00	0.00	0.00	0.05	0.00	0.40	0.00	1.30	0.00	0.01	0.00	0.00
29th	0.00		0.00	0.00	0.03	0.00	0.00	1.16	0.00	0.00	0.00	0.00
30th	0.00		0.00	0.00	0.01	0.00	0.00	2.00	0.00	0.04	0.00	0.00
31st	0.00		0.00		0.00		0.00	0.36		0.00		0.00

(Monthly Figures below)

Year's Rainfall Total:

45.77

SUM	2.94	2.32	2.12	0.89	3.15	3.25	6.82	10.05	3.93	2.09	3.42	4.79
MAX	0.82	1.65	1.30	0.42	1.20	1.54	2.30	2.20	1.50	0.35	1.35	1.20
MIN	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
AVG	0.09	0.08	0.07	0.03	0.10	0.11	0.22	0.32	0.13	0.07	0.11	0.15

"ND" or "-" = missing data / Shaded Cells = date samples were taken  
(Shellfish Management Areas 02 and 03)

# ANNUAL TABLE OF RAINFALL DATA

*SOURCE: Department of Natural Resources*

*Land, Water and Conservation Division*

*State Climatology Office (Myrtle Beach, SC / Station #2)*

2003	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
1st	--	0.01	0.03	0.00	0.00	0.27	0.00	0.00	0.00	0.00	0.00	0.00
2nd	0.50	0.00	0.90	0.00	0.00	0.00	0.16	0.00	0.00	0.00	0.00	0.00
3rd	0.10	0.00	0.00	0.00	0.12	0.00	0.28	0.00	0.00	0.00	0.06	--
4th	0.05	0.00	0.00	0.00	0.34	0.45	0.00	0.04	0.00	0.00	0.00	--
5th	0.00	0.35	0.07	0.00	0.00	0.65	0.00	0.10	0.50	0.00	0.00	--
6th	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
7th	0.00	0.74	0.82	0.35	0.67	1.40	0.00	0.05	0.00	0.00	0.00	0.00
8th	0.00	0.00	--	1.00	0.00	0.39	0.19	0.00	0.07	0.20	0.00	0.00
9th	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.10	0.00	0.23	0.00
10th	0.00	0.40	0.00	0.90	0.00	0.00	0.00	0.10	0.03	0.00	0.00	--
11th	0.00	0.10	0.00	0.35	0.00	0.00	0.00	0.67	0.00	0.00	0.00	1.05
12th	0.00	0.00	0.00	0.00	0.00	0.00	0.74	0.00	0.00	0.06	0.00	--
13th	0.00	0.00	0.00	0.00	0.00	0.00	2.16	0.01	0.00	0.00	0.00	0.00
14th	0.00	0.00	0.05	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.01	3.16
15th	0.00	0.00	--	0.00	0.26	0.00	0.45	0.35	0.00	0.00	0.00	--
16th	0.00	0.00	1.64	0.00	0.13	0.18	0.00	0.00	0.00	0.00	0.00	--
17th	0.13	0.51	0.00	0.00	0.40	0.73	0.00	0.00	0.00	0.00	0.00	--
18th	0.00	0.01	0.45	0.00	0.01	0.00	0.00	0.35	0.01	0.01	0.00	--
19th	0.00	0.00	0.00	0.00	0.01	0.67	0.80	0.27	0.43	0.00	1.11	--
20th	0.00	0.00	0.00	0.00	0.00	3.10	1.30	0.00	0.00	0.00	0.00	0.00
21st	--	0.00	2.35	0.00	0.00	0.78	0.00	0.00	0.00	0.00	0.00	--
22th	0.10	0.00	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
23rd	0.00	0.35	0.00	0.00	1.57	0.00	0.14	0.00	0.30	0.00	0.00	0.00
24th	0.00	0.00	0.00	0.00	0.16	0.00	2.26	0.00	0.05	0.00	0.00	0.18
25th	0.00	0.00	0.00	0.00	0.00	0.00	0.53	0.05	0.00	0.00	0.00	--
26th	0.00	0.00	0.00	1.76	--	0.00	0.00	0.00	0.00	0.00	0.00	--
27th	0.00	0.18	0.00	0.10	0.01	0.00	--	0.00	0.00	0.13	0.00	--
28th	0.00	0.30	0.00	0.00	0.13	0.00	0.05	0.00	0.00	0.10	0.13	--
29th	0.00		0.20	0.00	0.00	0.24	0.00	0.00	0.00	0.00	0.00	--
30th	0.08		0.00	0.00	0.14	0.00	0.00	0.00	0.00	0.00	0.00	0.00
31st	0.01		0.20		0.01		0.00	0.00		0.00		0.00

(Monthly Figures below)

Year's Rainfall Total:

46.93

SUM	0.97	2.95	6.72	4.47	3.96	8.86	9.09	1.99	1.49	0.50	1.54	4.39
MAX	0.50	0.74	2.35	1.76	1.57	3.10	2.26	0.67	0.50	0.20	1.11	3.16
MIN	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
AVG	0.03	0.11	0.23	0.15	0.13	0.30	0.30	0.06	0.05	0.02	0.05	0.29

"ND"or"--" = missing data / Shaded Cells = date samples were taken  
(Shellfish Management Areas 02 and 03)